

# Efficacy and safety assessment of rhamnolipids for use as a disease suppression agent in citrus trees infected with *Candidatus Liberibacter asiaticus*

Submitted by: Carolyn Slupsky, Professor, UC Davis.

## Phase 1: Preliminary experiments (*in vitro*)

### Research questions

- 1a. Can CLAs be cultured?
- 1b. What happens when rhamnolipids are applied to a culture of CLAs?
- 1c. Development of methods to detect rhamnolipids

### Experiments

#### 1a-b. Preparation and testing of *Candidatus Liberibacter asiaticus* (CLAs) culturing

Media that is capable of cultivating CLAs will be prepared in accordance with the methodology developed by our collaborators. Due to its limited capacity for aerobic respiration (Lin et al., 2011), CLAs will be cultured in an anaerobic growth chamber. Bacterial growth will be measured using one or more of the methodologies listed in **Table 1**. The methodology selected will depend on absolute quantities of CLAs that is able to be cultured and the associated environmental conditions (based on our collaborator's methods). The identity of CLAs will be confirmed by RT-qPCR (Bastianel et al., 2005), gram staining (Claus, 1992), and genomic sequencing (Zheng et al., 2015).

Proliferation and/or Survival	Bacterial growth can be measured by:  (1) viable counts of living bacteria <i>via</i> colony-forming unit (CFU) counts  (2) indirect total population count <i>via</i> turbidity (measured using a spectrophotometer)  (3) staining assay  (4) fluorescent based imaging	<b>Option 1</b> requires that CLAs be culturable on stationary media and is able to survive at low titers.  <b>Option 2</b> requires that CLAs be culturable in non-stationary media.  <b>Option 3</b> is ideal for determining the number of viable cells present in a cell suspension.  <b>Option 4</b> requires CLAs to be culturable for several passages (Caldwell et al., 1992).
-------------------------------	--	--

#### 1c. Development of methods to detect rhamnolipid

Rhamnolipids have been previously characterized by NMR using 1-dimensional (<sup>1</sup>H, <sup>13</sup>C, DEPT-135) and 2-dimensional (COSY, HMQC, HMBC) methods using deuterated methanol as the calibration standard; other methods include Thin-Layer Chromatography, LC-MS, and MS-MS (Sharma et al., 2007). Methods will be developed to detect the presence of this compound in the plant.

## Phase 2: Efficacy and validation experiments (greenhouse)

### *Research questions*

- 2a. What happens to citrus metabolism when rhamnolipids are injected into a citrus tree?
- 2b. What happens when rhamnolipids are injected into a citrus tree infected with CLAs?
- 2c. What happens to other bacteria in the rhizobiome and phytobiome when rhamnolipids are injected into a citrus tree +/- CLAs

### *Experiments*

#### 2a. Detecting rhamnolipids in citrus plant tissue

Greenhouse-grown citrus trees (n=12) from one variety (orange, lemon, OR mandarin) will be injected with rhamnolipids through the hypocotyl (stem area between the roots and canopy). Trees will be destructively sampled after 1d, 7d, 14d, and 21d post-exposure and leaves, stems, roots, and branches will be collected for analysis. Rhamnolipids and plant metabolites will be extracted and analyzed from plant tissue using methodology developed in our lab.

This experiment will answer the following questions: (1) At what concentration and for how long can rhamnolipids be detected in plant tissue; (2) Once injected, are rhamnolipids detectable in all parts of the plant (except fruit at this stage); and (3) Do rhamnolipids alter metabolism of healthy citrus?

#### 2b-c. Evaluating the *in planta* effect of rhamnolipids on CLAs-exposed citrus trees

One-year old greenhouse-grown trees (n=40) from one variety of citrus (orange, lemon, OR mandarin) will be grown at the contained research facility (CRF). Trees will be divided into 5 groups: (1) prevention (n=10, CLAs exposure, rhamnolipid treatment 1 day post-exposure); (2) early-stage disease (n=10, CLAs exposure followed by rhamnolipid treatment when infection confirmed by the metabolomics early detection method); (3) late-stage disease (n=10, CLAs exposure followed by rhamnolipid treatment when visual symptoms appear); (4) control (n=5, no CLAs exposure, rhamnolipids are applied at the start of the experiment); (5) true control (n=5, no CLAs exposure, no rhamnolipid treatment). Treatment plants will be exposed to CLAs *via* grafting with infected budwood or direct injection of the pathogen from culture. Control trees will receive grafts from uninfected budwood or direct injection of the vehicle. The visual symptoms of HLB used to classify disease progression will include: curled/deformed leaves, chlorosis, corked veins, mottling, twig dieback, thick/leathery leaves. Samples will be collected according to the schedule outlined in **Table 2**. Leaf samples will be collected for RT-qPCR (monthly), <sup>1</sup>H NMR-based metabolomics and/or MS/MS analysis for citrus metabolites and rhamnolipid detection respectively (monthly), and microbiome (every other month for the phytobiome) until plants fruit (approx. 12 months), after which they will be destructively sampled (leaves, hypocotyl, stems, roots, fruits, soil, flowers, bark).

This experiment will answer the following questions: (1) Do rhamnolipids prevent acquisition of CLAs; (2) At what stage of disease are rhamnolipids most effective (early (before symptom



2b-c. Evaluating the <i>in planta</i> effect of rhamnolipids on CLAs-exposed citrus trees													
Data Analysis													
<b>Manuscript Preparation</b>													
Phase 2													

Once Phase 1 is completed, we will work on defining the experiments for phase 3 of the study. We propose that the project be completed in 3 phases. Phase 1 will serve to replicate the results from Brazil in our hands. Phase 2 will begin at the same time to show that rhamnolipids are effective at killing CLAs *in planta*. Toward the end of phase 2, we will begin phase 3 to assess the safety in humans / animals / the environment. We envision that many manuscripts will be generated from these results. These studies will test: (1) the effectiveness of rhamnolipids in treating citrus infected by HLB, (2) the impact on the health and safety of humans or animals consuming fruit from a tree treated with rhamnolipids; (3) whether rhamnolipids change the rhizosphere, phyllosphere, and soil microorganisms of citrus trees; (4) whether rhamnolipids pose a threat to beneficial insects; and (5) potential impacts on ACP and their ability to carry the bacterium.

## References

- Bastianel, C., Garnier-Semancik, M., Renaudin, J., Bove, J.M., Eveillard, S. (2005). Diversity of “*Candidatus Liberibacter asiaticus*,” based on the *omp* gene sequence. *Appl Environ Microbiol*, 71(11), 6473-6478. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1287744/>
- Caldwell, D.E., Korber, D.R., Lawrence, J.R. (1992). Imaging of bacterial cells by fluorescence exclusion using scanning confocal laser microscopy. *J Micro Meth*, 15(4), p.249-261. <https://www.sciencedirect.com/science/article/pii/0167701292900456>
- Claus, D. (1992). A standardized Gram staining procedure. *World J Micro Biotech*, 8(4), p. 451-452. <https://link.springer.com/article/10.1007%2FBF01198764?LI=true>
- Iqbal, J., Siddiqui, R., Kazmi, S. U., & Khan, N. A. (2013). A simple assay to screen antimicrobial compounds potentiating the activity of current antibiotics. *BioMed Research International*, 2013.
- Lin, H., Lou, B., Glynn, J.M., Doddapaneni, H., Civerolo, E.L., Chen, C., Duan, Y., Zhou, L., Vahling, C.M. (2011). The complete genome sequence of ‘*Candidatus Liberibacter solanacearum*’, the bacterium associated with potato zebra chip disease. *PLoS*, 6(4), e19135. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3084294/#pone.0019135-Vahling1>

Sharma, A., Jansen, R., Nimtz, M., Johri, B.N., Wray, V. (2007) Rhamnolipids from the rhizosphere bacterium *Pseudomonas* sp. GRP3 that reduces damping-off disease in chilli and tomato nurseries. *J Nat Prod*, 70, 941-947. <https://pubs.acs.org/doi/abs/10.1021/np0700016>

Strober, W. (2001). Trypan blue exclusion test of cell viability. *Curr Protoc Immunol*, Appendix 3B, <https://www.ncbi.nlm.nih.gov/pubmed/18432654>

Zheng, Z., Sun, X., Deng, X., Chen, J. (2015) Whole-genome sequence of “*Candidatus Liberibacter asiaticus*” from Huanglongbing-affected citrus tree in central Florida. *Genome Annouce*, 3(2), e00169-15. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4395054/>